COVID-19 AND DEMENTIA



The below algorithm should be read in conjunction with official HSE guidance on www.hpsc.ie / www.hselibrary.ie

COVID-19 TESTING

Testing for COVID-19 can be **distressing and overwhelming** for many people with dementia.

- Explaining the testing procedure is important;
- This may be more difficult for people with advanced dementia to understand;
- Ensure all practicable steps are taken to enable understanding of the procedure and what it entails.

In Residential Care Facilities staff should be advised to have a **list of residents** who they feel will be unable to take part and may become verbally or physically distressed as a result.

Note: In some cases where clinical presentation suggests a diagnosis of COVID-19 and it is not possible to gain consent <u>or where it would be too</u> <u>stressful for the person with dementia</u>, the medical officer may decide not to refer for testing and in such cases a positive diagnosis would be assumed.

USE of PPE

PPE can be frightening for the person with dementia. They may not understand why the equipment is being worn; they may not recognize staff or pick-up facial and body cues to aid communication.

V

How you communicate becomes very important:

- Get the person's attention by touching their arm;
- Remain calm, use a reassuring, positive tone of voice;
- If possible establish clear YES-NO signals e.g. thumbs up for yes etc.
- Use other gestures as appropriate;
- If possible have a photo of yourself to hand;
- Use smiley stickers that can be discarded with the equipment.

